

On Integral Operators

Khalida Inayat Noor and Muhammad Aslam Noor

Department of Mathematics and Statistics Dalhousie University Halifax Nova Scotia

metadata, citation and similar papers at core.ac.uk

Submitted by George Leitmann

Received April 12, 1999

Let $f_n(z) = z/(1-z)^{n+1}$, $n \in N_o$, and $f_n^{(-1)}$ be defined such that $f_n * f_n^{(-1)} = \frac{z}{1-z}$, where $*$ denotes convolution (Hadamard product). Let f be analytic in the unit disc E . We define a new operator $I_n f = f_n^{(-1)} * f$ which is analogous to one defined by Ruscheweyh. Using this operator, the classes $M_{(n)}^*$ are defined. A function f , analytic in E , is in $M_{(n)}^*$ if and only if $I_n f$ is close-to-convex. The properties of $f \in M_{(n)}^*$ are discussed in some detail. It is shown that $M_{(n)}^* \subset M_{(n+1)}^*$ for $n \in N_o$ and for $n = 0, 1$, $M_{(n)}^*$ consists entirely of univalent functions. Closure properties of some integral operators defined on $M_{(n)}^*$ are also given. © 1999

Academic Press

Key Words: convolution; Ruscheweyh derivative; close-to-convex; univalent; integral operator.

1. INTRODUCTION

Let A denote the class of analytic functions defined in the unit disc $E = \{z : |z| < 1\}$ and normalized by $f(0) = 0$, $f'(0) = 1$. Let S , K , S^* , and C be the subclasses of A which contain, respectively, the univalent, close-to convex, starlike and convex functions.

Let $f \in A$. Denote by D^α : $A \rightarrow A$ the operator defined by

$$D^\alpha f(z) = \frac{z}{(1-z)^{\alpha+1}} * f(z), \quad \alpha > -1,$$

which implies that

$$D^n f(z) = \frac{z(z^{n-1}f(z))^{(n)}}{n!}$$

for $n \in N \cup \{0\} = N_o$ and the operator $(*)$ stands for the Hadamard product or convolution. We note that $D^o f(z) = f(z)$ and $D'f(z) = zf'(z)$. The operator $D^n f$ is called the Ruscheweyh derivative of n th order of f . Several classes of analytic functions, defined by using this operator, have been studied, see [1, 2, 8, 10] and others.

Analogous to $D^n f$, we here define an integral operator $I_n: A \rightarrow A$ as follows.

Let $f_n(z) = z/(1-z)^{n+1}$, $n \in N_o$, and let $f_n^{(-1)}$ be defined such that

$$f_n(z) * f_n^{(-1)}(z) = \frac{z}{1-z}. \quad (1.1)$$

Then

$$I_n f = f_n^{(-1)} * f = \left[\frac{z}{(1-z)^{n+1}} \right]^{(-1)} * f. \quad (1.2)$$

Using (1.1), (1.2), and a well-known identity for $D^n f$, we have

$$(n+1)I_n f - nI_{n+1} f = z(I_{n+1} f)'. \quad (1.3)$$

Using hypergeometric functions ${}_2F_1$, we can write (1.2) as

$$I_n f = [z {}_2F_1(1, 1; n+1, z)] * f.$$

We note that $I_0 f' = zf$ and $I_1 f = f$. We now define

DEFINITION 1.1. Let $f \in A$. Then $f \in N_{(n)}^*$, $n \in N_o$ if and only if $I_n f \in S^*$ for $z \in E$. It is clear that $N_{(o)}^* = C$ and $N_{(1)}^* = S^*$. The classes $N_{(n)}^*$ have been studied in [7] and it is shown that $N_{(n)}^* \subset N_{(n+1)}^*$ for each $n \in N_o$. We here extend the classes $N_{(n)}^*$ as

DEFINITION 1.2. Let $f \in A$. The $f \in M_{(n)}^*$ for $n \in N_o$ if and only if there exists $g \in N_{(n)}^*$ such that, for $z \in E$,

$$\operatorname{Re} \left\{ \frac{z(I_n f(z))'}{I_n g(z)} \right\} > 0.$$

We note that $M_{(1)}^* = K$ and $M_{(o)}^* = C^*$, where C^* is the class of quasi-convex functions first introduced and studied in [6].

2. PRELIMINARY RESULTS

In order to develop some results for the classes $M_{(n)}^*$, we need

LEMMA 2.1 [4]. Let ω be analytic in E . If $|\omega|$ assumes its maximum value on the circle $z = r$ at a point z_o , then $z_o \omega'(z_o) = k \omega(z_o)$ where $k \geq 1$.

LEMMA 2.2 [9]. Let $\phi \in C$ and $g \in S^*$ in E . Then for F analytic in E with $F(0) = 1$, $\phi^* F g / \phi^* g$ is contained in the convex hull of $F(E)$.

LEMMA 2.3 [5]. Let $u = u_1 + iu_2$ and $v = v_1 + iv_2$ and let $\psi(u, v)$ be a complex-valued function satisfying the conditions:

- (i) $\psi(u, v)$ is continuous in $D \subset \mathbb{C}^2$,
- (ii) $(1, 0) \in D$ and $\psi(1, 0) > 0$,
- (iii) $\operatorname{Re} \psi(iu_2, v_1) \leq 0$ whenever $(iu_2, v_1) \in D$ and $v_1 \leq -\frac{1}{2}(1 + u_2^2)$.

If $h(z) = 1 + \sum_{k=2}^{\infty} c_k z^k$ is a function analytic in E such that $(h(z), zh'(z)) \in D$ and $\operatorname{Re} \psi(h(z), zh'(z)) > 0$ for $z \in E$, then $\operatorname{Re} h(z) > 0$ in E .

LEMMA 2.4. Let p be analytic in E with $p(0) = 1$ and $\operatorname{Re} p(z) > 0$, $z \in E$. Then, for $|z| = r$, $z \in E$,

- (i) $\frac{1-r}{1+r} \leq \operatorname{Re} p(z) \leq |p(z)| \leq \frac{1+r}{1-r}$
- (ii) $|p'(z)| \leq 2 \operatorname{Re} p(z) / (1 - r^2)$.

3. MAIN RESULTS

The following is an inclusion result for the family $M_{(n)}^*$.

THEOREM 3.1. $M_{(n)}^* \subset M_{(n+1)}^*$ for each $n \in N_o$.

In particular $M_{(o)}^* \subset M_{(1)}^*$ which implies $C^\alpha \subset K$. This result is known [6].

Proof. Let $f \in M_{(n)}^*$. Then, for $z \in E$,

$$\operatorname{Re} \left\{ \frac{z(I_n f(z))'}{I_n g(z)} \right\} > 0, \quad \text{for some } g \in N_{(n)}^*.$$

Define $\omega(z)$ in E such that

$$\frac{z(I_{n+1}(z))^*}{I_{n+1}(z)} = \frac{1 - \omega(z)}{1 + \omega(z)}, \quad (3.1)$$

where $\omega(0) = 0$ and $\omega(z) \neq -1$. We show that $|\omega(z)| < 1$. From (3.1) and the identity (1.3), we have

$$(n+1) \frac{z(I_n f(z))'}{I_{n+1} g(z)} = \frac{z(I_{n+1} g(z))'}{I_z g(z)} \left[\frac{1 - \omega(z)}{1 + \omega(z)} \right] + \frac{I_{n+1} g(z)}{I_n g(z)} \left\{ \frac{-2z\omega'(z)}{[1 + \omega(z)]^2} + n \left[\frac{1 - \omega(z)}{1 + \omega(z)} \right] \right\}. \quad (3.2)$$

We now apply (1.3) for the function g and since $N_{(n)}^* \subset N_{(n+1)}^*$, there exists an analytic function $\omega_1(z)$ with $\omega_1(0) = 0$ and $|\omega_1(z)| < 1$ such that

$$\frac{I_n g(z)}{I_{n+1} g(z)} = \frac{1 - \omega_1(z)}{1 + \omega_1(z)}. \quad (3.3)$$

Thus, from (3.2) and (3.3), we have

$$\frac{z(I_n f(z))'}{I_n g(z)} = \frac{1 - \omega(g)}{1 + \omega(g)} + \frac{1}{n+1} \left(\frac{1 + \omega_1(z)}{1 - \omega_1(z)} \right) \left(\frac{2z\omega'(z)}{[1 + \omega(z)]^2} \right). \quad (3.4)$$

Suppose now that, for $z \in E$,

$$\max_{|z| \leq |z_o|} |\omega(z)| = |\omega(z_o)| = 1 \quad (\omega(z_o) \neq -1).$$

Then, from Lemma 2.1, it follows that

$$z_o \omega'(z_o) = k \omega(z_o), \quad \text{where } k \geq 1.$$

Setting $\omega(z_o) = e^{i\theta}$ and $\omega_1(z_o) = re^{i\phi}$ in (3.4) gives us

$$\begin{aligned} & \operatorname{Re} \left\{ \frac{z_o(I_n f(z_o))'}{I_n g(z_o)} \right\} \\ &= \operatorname{Re} \left\{ \frac{1}{n+1} \cdot \frac{-2k(e^{i\theta} + e^{-i\theta} + 2)(1 + r^2 + 2r \cos \phi)}{|1 + re^{i\phi}|^2 |1 + e^{i\theta}|^2} \right\} \\ &= \frac{-4k}{n+1} \left\{ \frac{(\cos \theta + 1)(1 + r^2 + 2r \cos \phi)}{|1 + re^{i\phi}|^2 |1 + e^{i\theta}|^2} \right\}. \end{aligned}$$

Hence, if $\phi = \frac{\pi}{2}$,

$$\operatorname{Re} \left\{ \frac{z_o (I_n f(z_o))'}{I_n g(z_o)} \right\} < 0, \quad \text{where } g \in N_{(n)}^* \text{ and } k \geq 1.$$

This contradicts our hypothesis that $f \in M_n^*$. Thus $|\omega(z)| < 1$ and so, from (3.1), $f \in M_{(n+1)}^*$. Since it is known that $C^* \subset K$ and $f \in K$ is univalent, we note that $f \in M_{(n)}^*$ is univalent in E for $n = 0, 1$. ■

Remark 3.1. From (1.2) and Definition 1.2, it follows that

$$f \in M_{(n)}^* \text{ if and only if } I_n f \in K. \quad (3.5)$$

THEOREM 3.2. Let $f \in M_{(n)}^*$ and be given by $f(z) = z + \sum_{k=2}^{\infty} a_k z^k$. Then

$$|a_k| \leq \frac{(k+n-1)!}{n!(k-1)!}, \quad k \geq 2, n \in N_o.$$

This result is sharp.

Proof. $f \in M_{(n)}^*$ implies that $F = I_n f \in K$. Therefore $F = f_n^{(-1)} * f$ and so $f_n * F = \frac{z}{1-z} * f$.

Let $F(z) = z + \sum_{k=2}^{\infty} b_k z^k$. Then

$$|a_k| = \frac{(n+k-1)!}{n!k!} |b_k| \leq \frac{(n+k-1)!}{n!(k-1)!} \quad (|b_k| \leq \text{since } F \in K),$$

and this gives us the required result. The sharpness follows if we take

$$I_n f_0(z) = \frac{z}{(1-z)^2}; \quad f_0^{(z)} = z + \sum_{k=2}^{\infty} \frac{(k+n-1)!}{n!(k-1)!} z^k.$$

■

Special Cases. (i) $f \in M_{(0)}^* \equiv C^*$ implies $|a_k| \leq 1$ for all $k \geq 2$. This result is known [6].

(ii) For $f \in M_{(1)}^* \equiv K$, $|a_k| \leq k$ for all $k \geq 2$, see [3].

(iii) Let $n \in N_o$ and $k = 2$. Then

$$|a_2| \leq (n+1).$$

Using Theorem 3.2 and the fact that $f \in M_{(n)}^*$ is univalent for $n = 0, 1$, we can immediately prove the following covering result.

THEOREM 3.3. Let $f \in M_{(n)}^*$, $n = 0, 1$. If B is the boundary of the image of E under f , then every point of B is at distance at least $\frac{1}{n+3}$ from the origin.

THEOREM 3.4. Let $f \in M_{(n+1)}^*$. Then $f \in M_{(n)}^*$ for $|z| < r_n = (1+n)/(2+\sqrt{n^2+3})$. This result is sharp.

Proof. Since $f \in M_{(n+1)}^*$, we can write

$$\frac{z(I_{n+1}f(z))'}{I_{n+1}g(z)} = H(z), \quad \operatorname{Re} H(z) > 0,$$

$$z \in E \quad \text{for some } g \in N_{(n+1)}^*.$$

Identity (1.3) together with some computations yields

$$\left\{ \frac{z(I_n f(z))'}{I_n g(z)} \right\} = \left\{ H(z) + \frac{zH'(z)}{h(z) + n} \right\},$$

where

$$h(z) = \frac{z(I_{n+1}g(z))'}{I_{n+1}g(z)} \quad \text{and} \quad \operatorname{Re} h(z) > 0 \text{ in } E.$$

Therefore

$$\begin{aligned} \operatorname{Re} \left\{ \frac{z(I_n f(z))'}{I_n g(z)} \right\} &= \operatorname{Re} \left\{ H(z) + \frac{zH'(z)}{h(z) + n} \right\} \\ &\geq \operatorname{Re} H(z) \left\{ 1 - \frac{2r}{1-r^2} \cdot \frac{1}{(1-r)/(1+r) + n} \right\} \\ &= \operatorname{Re} H(z) \left\{ \frac{(1+n) - 4r + (1-n)r^2}{(1-r)^2 + n(1-r^2)} \right\}, \end{aligned}$$

where we have used Lemma 2.4.

The right-hand side of the above inequality is positive for $|z| = r < r_n[(1+n)/(2+\sqrt{n^2+3})]$.

By taking $I_n f(z) = z(1-z)^2$ we see that the radius r_n is exact. ■

To prove our next result, we need

LEMMA 3.1. Let $G \in N_{(n)}^*$ and let, for $0 < \lambda \leq 1$, $g \in A$ be defined by

$$g(z) = \frac{1}{\lambda} z^{1-1/\lambda} \int_0^z t^{1/\lambda-2} G(t) dt. \quad (3.6)$$

Then

$$\operatorname{Re} \left\{ \frac{z(I_n g(z))'}{I_n g(z)} \right\} > \alpha \quad (0 < \alpha < 1),$$

where

$$\alpha = \frac{2\lambda}{\left[(\lambda - 2) + \sqrt{9\lambda^2 - 4\lambda + 4} \right]}. \quad (3.7)$$

Proof. Set $z(I_n g(z))'/I_n g(z) = (1 - \alpha)p(z) + \alpha$, $p(z) = 1 + c_1 z + c_2 z^2 + \dots$.

Then, from (3.6), we have

$$\frac{z(I_n g(z))'}{I_n g(z)} = (1 - \alpha)p(z) + \alpha + \frac{\lambda(1 - \alpha)zp'(z)}{1 - \lambda + \alpha\lambda + \lambda(1 - \alpha)p(z)}.$$

We now form the functional $\psi(u, v)$ by taking $u = p$ and $v = zp'$. So

$$\psi(u, v) = (1 - \alpha)u + \alpha + \frac{\lambda(1 - \alpha)v}{1 - \lambda + \alpha\lambda + \lambda(1 - \alpha)u}.$$

For

$$D = \mathbb{C} \setminus \left\{ -\frac{1 - \lambda + \alpha\lambda}{\lambda(1 - \alpha)} \right\} \times \mathbb{C},$$

the conditions (i) and (ii) of Lemma 2.3 are clearly satisfied. We proceed to verify condition (iii).

$$\begin{aligned} \operatorname{Re} \psi(iu_2, v_1) &= \alpha + \operatorname{Re} \frac{\lambda(1 - \alpha)v_1}{1 - \lambda + \alpha\lambda + i\lambda(1 - \alpha)u_2} \\ &= \frac{\alpha + \lambda(1 - \alpha)(1 - \lambda + \alpha\lambda)v_1}{(1 - \lambda + \alpha\lambda)^2 + \lambda^2(1 - \alpha)^2 u_2^2} \\ &\leq \alpha - \left(\frac{1}{2} \right) \frac{\lambda(1 - \alpha)(1 - \lambda + \alpha\lambda)(1 + u_2^2)}{(1 - \lambda + \alpha\lambda)^2 + \lambda^2(1 - \alpha)^2 u_2^2} \\ &= \frac{A + Bu_2^2}{2C}, \end{aligned}$$

where

$$\begin{aligned} A &= 2\alpha(1 - \lambda + \alpha\lambda)^2 - \lambda(1 - \alpha)(1 - \lambda + \alpha\lambda), \\ B &= 2\alpha\lambda^2(1 - \alpha)^2 - \lambda(1 - \alpha)(1 - \lambda + \alpha\lambda), \\ C &= (1 - \lambda + \alpha\lambda)^2 + \lambda^2 u_2^2(1 - \alpha)^2 > 0. \end{aligned}$$

$\operatorname{Re} \psi(iu_2, v_1) \leq 0$ if and only if $A \leq 0$ and $B \leq 0$. From $A \leq 0$, we obtain α as given by (3.7) and $B \leq 0$ gives us $0 < \alpha < 1$. ■

THEOREM 3.5. Let $F \in M_{(n)}^*$ and let f be defined by

$$f(z) = \frac{1}{\lambda} z^{1-1/\lambda} \int_0^z t^{(1/\lambda)-2} F(t) dt \quad (3.8)$$

for $0 < \lambda \leq 1$. Then there exists a function g with

$$\operatorname{Re} \left\{ \frac{z(I_n g(z))'}{I_n g(z)} \right\} > \alpha$$

such that

$$\operatorname{Re} \left\{ \frac{z(I_n f(z))'}{I_n g(z)} \right\} > \beta, \quad \text{for } z \in E,$$

where α is given by (3.7) and β ($0 \leq \beta < 1$) is defined by (3.14).

Proof. Let $G \in N_{(n)}^*$ and let, for $0 < \lambda \leq 1$, g be given by (3.6). Then we can write

$$I_n G(z) = (1 - \lambda)I_n g(z) + \lambda z(I_n g(z))',$$

and

$$I_n F(z) = (1 - \lambda)I_n f(z) + \lambda z(I_n f(z))'.$$

Set

$$\frac{z(I_n f(z))'}{I_n g(z)} = (1 - \beta)h(z) + \beta, \quad h(0) = 1. \quad (3.9)$$

We show that $\operatorname{Re} h(z) > 0$ in E .

From the above relations, we have

$$\frac{z(I_n F(z))'}{I_n g(z)} = \left\{ \frac{(1-\lambda)z(I_n f(z))'}{I_n g(z)} + \frac{\lambda z[z(I_n f(z))']'}{I_n g(z)} \right\} / \left[(1-\lambda) + \frac{\lambda z(I_n g(z))'}{I_n g(z)} \right]. \quad (3.10)$$

From Lemma 3.1,

$$\frac{z(I_n g(z))'}{I_n g(z)} = (1-\alpha)p(z) + \alpha, \quad \operatorname{Re} p(z) > 0 \quad \text{in } E. \quad (3.11)$$

From (3.9) and (3.11) we obtain

$$\begin{aligned} \frac{z[z(I_n f(z))']'}{I_n g(z)} &= \{(1-\alpha)p(z) + \alpha\}\{(1-\beta)h(z) + \beta\} \\ &\quad + (1-\beta)zh'(z). \end{aligned} \quad (3.12)$$

Using (3.9), (3.11), and (3.12) in (3.10), we have

$$\frac{z(I_n F(z))'}{I_n G(z)} = \beta + (1-\beta)h(z) + \frac{\lambda(1-\beta)zh'(z)}{(1-\lambda) + \lambda[(1-\alpha)p(z) + \alpha]}. \quad (3.13)$$

We now define $\psi(u, v)$ by taking $u = h(z)$ and $v = zh'(z)$ in (3.13) as

$$\psi(u, v) = \beta + (1-\beta)u + \frac{\lambda(1-\beta)v}{\lambda(1-\alpha)p - \lambda(1-\alpha) + 1}.$$

It is clear that $\psi(u, v)$ satisfies conditions (i) and (ii) of Lemma 2.3. To verify condition (iii), we note that

$$\operatorname{Re} \psi(iu_2, v_1) = \beta + \frac{\lambda(1-\beta)v_1\{\lambda(1-\alpha)p_1 - \lambda(1-\alpha) + 1\}}{\{\lambda(1-\alpha)p_1 - \lambda(1-\alpha) + 1\}^2 + \lambda^2(1-\alpha)^2 p_2^2},$$

where $p(z) = p_1 + ip_2$, p_1, p_2 being functions of x and y and $\operatorname{Re} p = p_1 > 0$ in E . By putting $v_1 \leq -\frac{1}{2}(1 + u_2^2)$, we obtain

$$\begin{aligned} & \operatorname{Re} \psi(iu_2, v_1) \\ & \leq \beta - \frac{\lambda(1 - \beta)(1 + u_2^2)[\lambda(1 - \alpha)p_1 - \lambda(1 - \alpha) + 1]}{2[\{\lambda(1 - \alpha)p_1 - \lambda(1 - \alpha) + 1\}^2 + \lambda^2(1 - \alpha)^2 p_2^2]} \\ & = \frac{A + Bu_2^2}{2C}, \end{aligned}$$

where

$$\begin{aligned} C &= \{\lambda(1 - \alpha)p_1 - \lambda(1 - \alpha) + 1\}^2 + \lambda^2(1 - \alpha)^2 p_2^2 > 0, \\ A &= 2\beta\{\lambda(1 - \alpha)p_1 - \lambda(1 - \alpha) + 1\}^2 + \lambda^2(1 - \alpha)^2 p_2^2 \\ &\quad - \lambda(1 - \beta)[\lambda(1 - \alpha)p_1 - \lambda(1 - \alpha) + 1], \end{aligned}$$

and

$$B = -\lambda(1 - \beta)[\lambda(1 - \alpha)p_1 - \lambda(1 - \alpha) + 1].$$

We note that $\operatorname{Re} \psi(iu_2, v_1) \leq 0$ if and only if $A \leq 0$ and $B \leq 0$. From $A \leq 0$, we obtain $\beta \leq \beta_A$, where

$$\beta_A = \frac{\lambda^2(1 - \alpha)^2 p_2^2 + \lambda[\lambda(1 - \alpha)p_1 - \lambda(1 - \alpha) + 1]}{2[\{\lambda(1 - \alpha)p_1 - \lambda(1 - \alpha) + 1\}^2 + \lambda^2(1 - \alpha)^2 p_2^2] + \{\lambda(1 - \alpha)p_1 - \lambda(1 - \alpha) + 1\}}. \quad (3.14)$$

Also, from $B \leq 0$, we have $\beta_A < 1$ and the condition (iii) is satisfied to give $\operatorname{Re} h(z) > 0$ for $z \in E$. This completes the proof. ■

Let, for $f \in A$ and $n = (\frac{1}{\lambda} - 1)$ in (3.8),

$$\begin{aligned} f(z) &= L(F) = \frac{n+1}{z^n} \int_0^z t^{n-1} F(t) dt, \\ L(F) &= \left(z \sum_{k=0}^{\infty} \frac{n+1}{n+k+1} z^k \right) * F = \left(z \sum_{k=0}^{\infty} \frac{(n+1)_k (1)_k}{(n+2)_k k!} \right) * F \\ &= [z {}_2F_1(1, n+1; n+2, z)] * F \\ &= \frac{z}{(1-z)^{n+1}} * \left[\frac{z}{(1-z)^{n+2}} \right]^{(-1)} * F \\ &= f_n * f_{n+1}^{(-1)} * F. \end{aligned} \quad (3.15)$$

This implies that

$$I_n L(F) = I_{n+1} F.$$

Thus we have

THEOREM 3.6. *Let $F \in M_{(n+1)}^*$. Then $f = L(F) \in M_{(n)}^*$ for each $n \in N_0$ and $f = L(F)$ is defined by (3.15).*

Next we prove that the classes $M_{(n)}^*$ are closed under convolution with convex functions. Many results on $M_{(n)}^*$ can be deduced as applications of this result.

THEOREM 3.7. *Let $\phi \in C$ and $f \in M_{(n)}^*$. The $f * \phi \in M_{(n)}^*$.*

Proof. Let

$$\begin{aligned} I_n(f * \phi) &= f_n^{(-1)} * (f * \phi) \\ &= (f_n^{(-1)} * f) * \phi \\ &= (I_n f) * \phi. \end{aligned}$$

Since $I_n f \in K$ and $\phi \in C$, it follows that $I_n f * \phi \in K$, see [9]. Consequently $f * \phi \in M_{(n)}^*$. ■

We can apply Theorem 3.7 to note that the classes $M_{(n)}^*$ are invariant under the following operators.

- (i) $f_1(z) = \int_0^z \frac{f(t)}{t} dt$
- (ii) $f_2(z) = \int_0^z \frac{f(t) - f(xt)}{t - xt} dt, |x| \leq 1, x \neq 1$
- (iii) $f_3(z) = (1 + c)/z^c \int_0^z t^{c-1} f(t) dt, \operatorname{Re} c > 0.$

The proof follows immediately since we can write

$$f_i = f * \phi_i, \quad i = 1, 2, 3,$$

with

$$\begin{aligned} \phi_1(z) &= \sum_{k=1}^{\infty} \frac{z^k}{k} = -\log(1 - z), \\ \phi_2(z) &= \sum_{k=1}^{\infty} \frac{1 - x^k}{k(1 - x)} z^k = \frac{1}{1 - x} \log \frac{1 - xz}{1 - z} \quad (|x| = 1, x \neq 1), \\ \phi_3(z) &= \sum_{k=1}^{\infty} \frac{1 + c}{k + c} z^k, \quad \operatorname{Re} c > 0, \end{aligned}$$

and ϕ_i is convex for each $i = 1, 2, 3$.

REFERENCES

1. O. Ahuja, Integral operators of certain univalent functions, *Internat. J. Math. Math. Sci.* **8** (1985), 653–662.
2. H. S. Al-Amiri and T. S. Fernando, On close-to-convex functions of complex order, *Internat. J. Math. Math. Sci.* **13** (1990), 321–330.
3. A. W. Goodman, “Univalent Functions,” Vols. 1, 2, Polygonal Publishing House, Washington, NJ, 1983.
4. I. Jack, Functions starlike and convex of order α , *J. London Math. Soc.* **3** (1971), 469–474.
5. S. Miller, Differential inequalities and Caratheodory functions, *Bull. Amer. Math. Soc.* **81** (1975), 79–81.
6. K. I. Noor and D. K. Thomas, On quasi-convex univalent functions, *Internat. J. Math. Math. Sci.* **3** (1980), 225–266.
7. K. I. Noor, On new classes of integral operators, *J. Nat. Geometry* **16** (1999), 71–80.
8. K. I. Noor, On certain classes of close-to-convex functions, *Internat. J. Math. Math. Sci.* **16** (1993), 329–336.
9. S. Ruscheweyh and T. Shiel-Small, Hadamard products of schlicht functions and the Polya–Schoenberg conjecture, *Comment. Math. Helv.* **48** (1973), 119–135.
10. R. Singh and S. Singh, Integrals of certain univalent functions, *Proc. Amer. Math. Soc.* **77** (1979), 336–340.